

HOW TO USE



*and make the most out of your cutaneous
T-cell lymphoma (CTCL) treatment*

TARGRETIN capsules should not be used by women who are pregnant, believe they may be pregnant, or are planning to become pregnant.

Please see Important Safety Information, including Pregnancy Boxed Warning, located on the inside back panel, and Patient Instructions for Use located within the accompanying full Prescribing Information. Activate your card at Targretin.com.*

*This offer is only valid for patients with commercial insurance. Commercially insured patients whose insurance does not cover TARGRETIN will pay more. These savings offers are not valid for any person eligible for reimbursement of prescriptions, in whole or in part, by any federal, state, or other governmental programs, including, but not limited to, Medicare, (including Medicare Advantage and Part A, B and D Plans), Medicaid, TRICARE, Veterans Administration or Department of Defense health coverage, CHAMPUS, the Puerto Rico Government Health Insurance Plan, or any other federal or state health care programs. Void where prohibited by law, taxed or restricted. Cannot be redeemed at government-subsidized clinics. Ortho Dermatologics reserves the right to rescind, revoke, or amend this offer without notice. **See inside panel for continued eligibility terms and conditions.**

THE BASICS ABOUT TARGRETIN CAPSULES

TARGRETIN® (bexarotene) capsules are a prescription medicine used to treat the skin problems that happen with a type of cancer called cutaneous T-cell lymphoma, or CTCL. It is used when treatment with at least one other type of medicine by mouth or injection did not work or stopped working. It is not known if TARGRETIN is safe and effective in children.

THERE ARE SOME IMPORTANT THINGS TO KNOW BEFORE STARTING TARGRETIN CAPSULES TREATMENT:

- **TARGRETIN capsules can cause major damage to a fetus.**

If you are capable of becoming pregnant, you should take contraceptive measures to avoid pregnancy before, during and after treatment. It is recommended that 2 reliable forms of contraception be used together.

- TARGRETIN capsules can greatly increase blood levels of lipids (triglycerides and cholesterol), and these levels must be monitored and, if elevated, treated.
- TARGRETIN capsules can cause an underactive thyroid, and periodic blood tests will be needed to detect this. Medication to control the condition may be necessary.
- DO NOT take more than the recommended daily dietary allowance of vitamin A (4000 to 5000 International Units).
- Minimize exposure to sunlight and artificial ultraviolet light (sunlamps, tanning beds, etc.).

***Terms and Conditions, continued from front cover:**

You agree not to seek reimbursement for all or any part of the benefit received through this offer and are responsible for making any required reports of your use of this offer to any insurer or other third party who pays any part of the prescription filled. You must be 18 years of age or older to redeem this offer for yourself or a minor. You must present this coupon along with your prescription to participate in this program. You must activate this coupon before using by calling 1-888-201-1385 or visiting Targretin.com. Reimbursement limitations apply. Patient is responsible for all additional costs and expenses after reimbursement limits are reached. This coupon and offer are not health insurance.

Please see Important Safety Information including Pregnancy Boxed Warning, located on the inside back panel, and Patient Instructions for Use located within the accompanying full Prescribing Information.

HOW TO TAKE TARGRETIN (bexarotene) CAPSULES, 75 MG

TARGRETIN capsules should be taken once a day with a meal. Your doctor will tell you how many you need to take. Take them all at once, swallowing each capsule whole. If you miss a dose, take it as soon as possible. If it's almost time for your next dose, skip the missed dose.

HOW LONG BEFORE YOU CAN EXPECT YOUR CTCL TO IMPROVE ON TARGRETIN CAPSULE TREATMENT?

It is important to use the medication as directed by your physician as a response may take several weeks to several months. Your doctor should determine how long you should be taking TARGRETIN capsules.

*You should tell your doctor or nurse about any
current medications or health conditions.*

Please see Important Safety Information including Pregnancy Boxed Warning, located on the inside back panel, and Patient Instructions for Use located within the accompanying full Prescribing Information.

INDICATION

Targretin® (bexarotene) capsules are a prescription medicine used to treat the skin problems that happen with a type of cancer called cutaneous T-cell lymphoma, or CTCL. It is used when treatment with at least one other type of medicine by mouth or injection did not work or stopped working. It is not known if Targretin is safe and effective in children.

IMPORTANT SAFETY INFORMATION INCLUDING PREGNANCY BOXED WARNING

What are the most important things I should know about Targretin capsules?

Targretin can cause serious side effects, including major birth defects to an unborn baby, if taken during pregnancy.

- **You should avoid becoming pregnant during treatment with Targretin.**
- **Do not take Targretin if you are pregnant or plan to become pregnant.** Call your healthcare provider if you become pregnant or think you are pregnant while taking Targretin, and for one month after stopping treatment.
- **You must use two effective forms of birth control together starting one month before you begin treatment with Targretin, during treatment, and for one month after stopping Targretin.** Talk to your doctor about what forms of birth control are right for you.
- **You must use a condom if you are male and your partner is pregnant or capable of becoming pregnant.**

Do not take Targretin capsules if you are allergic to this medicine.

Before taking Targretin, tell your healthcare provider about all medical conditions, including if you:

- have or have had problems with your pancreas, including pancreatitis
- have or have had gallbladder problems
- have or have had liver problems
- have diabetes
- have high levels of fats (lipids) called cholesterol or triglycerides in your blood
- have cataracts or a history of cataracts
- have thyroid problems
- drink alcohol
- are breastfeeding or plan to breastfeed, or are pregnant or plan to become pregnant (see **“What are the Most Important Things I Should Know about Targretin?”**)
- are taking any other medications, including over-the-counter medicines and supplements

While taking Targretin You Should:

- Limit your exposure to the sun or artificial types of sunlight such as tanning beds and sunlamps
- Limit the amount of Vitamin A that you take

What are the possible side effects of Targretin?

- See **“What are the Most Important Things I Should Know about Targretin?”**
- **Increased levels of fats (lipids) called cholesterol or triglycerides in your blood are common with Targretin and can be serious.** It may require a dose adjustment, or interrupting or stopping treatment.
- **Inflammation of the pancreas (acute pancreatitis)** can sometimes be caused by Targretin and can lead to death. Ask your healthcare provider about factors that can increase the likelihood of developing pancreatitis.
 - **Tell your healthcare provider right away if you develop signs of pancreatitis,** such as nausea that doesn't go away, vomiting, back or stomach-area pain.
- **Liver problems,** including liver failure, can lead to death.
- **Thyroid problems** (hypothyroidism) are common with Targretin and may require treatment.
- **Low white blood cell count** is common with Targretin and sometimes severe.
- **New or worse cataracts** – tell your healthcare provider about any vision changes.
- **Risk of low blood sugar in people with diabetes** – talk to your doctor about your diabetes medicines to avoid drug interactions that can occur with Targretin.

The most common side effects of Targretin include headache, muscle weakness, rash, nausea, infection, stomach-area pain, swelling of hands, arms, feet or legs, and dry skin.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying full Prescribing Information including pregnancy Boxed Warning and Patient Information.



TARGRETIN (bexarotene) CAPSULES are a medicine used to treat the skin problems arising from a disease called CTCL after at least one other type of systemic therapy did not work or stopped working.

Before starting treatment with TARGRETIN capsules, make sure you tell your doctor about all your medical conditions and any medications you are currently taking.

Please see Important Safety Information including Pregnancy Boxed Warning, located on the inside back panel, and Patient Instructions for Use located within the accompanying full Prescribing Information.